

Finite Fields Appl. 35(2015), 86–91.

## PROOF OF A CONJECTURAL SUPERCONGRUENCE

XIANG-ZI MENG AND ZHI-WEI SUN\*

ABSTRACT. Let  $m > 2$  and  $q > 0$  be integers with  $m$  even or  $q$  odd. We show the supercongruence

$$\sum_{k=0}^{p-1} (-1)^{km} \binom{p/m - q}{k}^m \equiv 0 \pmod{p^3}$$

for any prime  $p > mq$ . This confirms a conjecture of Sun.

### 1. INTRODUCTION

Let  $p$  be a prime. A  $p$ -adic congruence is called a *supercongruence* if it happens to hold modulo higher powers of  $p$ . A classical result due to J. Wolstenholme (cf. [7] or [2]) states that if  $p > 3$  then

$$\sum_{k=1}^{p-1} \frac{1}{k} \equiv 0 \pmod{p^2} \quad \text{and} \quad \binom{2p-1}{p-1} \equiv 1 \pmod{p^3}.$$

For some recent supercongruences modulo  $p^2$  involving products of two binomial coefficients, one may consult [4] and [1].

Recently, Z.-W. Sun [5] established some new supercongruences modulo prime powers motivated by the well-known formula

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e.$$

For example, he obtained the following theorem.

**Theorem 1.1.** *For any prime  $p > 3$ , we have*

$$\sum_{k=0}^{p-1} \binom{-1/(p+1)}{k}^{p+1} \equiv 0 \pmod{p^5}$$

and

$$\sum_{k=0}^{p-1} \binom{1/(p-1)}{k}^{p-1} \equiv \frac{2}{3} p^4 B_{p-3} \pmod{p^5},$$

where  $B_0, B_1, B_2, \dots$  are the well-known Bernoulli numbers.

---

*Key words and phrases.*  $p$ -adic congruences, binomial coefficients.

2010 *Mathematics Subject Classification.* Primary 11A07; Secondary 05A10, 11B65.

\*Supported by the National Natural Science Foundation (grant 11171140) of China.

In this paper we show the following result conjectured by Sun [5].

**Theorem 1.2.** *Let  $m > 2$  and  $q > 0$  be integers with  $m$  even or  $q$  odd. Then, for any prime  $p > mq$ , we have the supercongruence*

$$\sum_{k=0}^{p-1} (-1)^{km} \binom{p/m - q}{k}^m \equiv 0 \pmod{p^3}. \quad (1.1)$$

We are going to provide few lemmas in the next section, and then show Theorem 1.2 in Section 3.

## 2. SOME LEMMAS

**Lemma 2.1.** *For any positive integer  $n$ , we have*

$$\sum_{k=0}^n \binom{n}{k} (-1)^k k^m = 0 \quad \text{for all } m = 0, 1, \dots, n-1.$$

*Remark 2.1.* This is a well-known result, see, e.g., [6, pp. 125-126].

**Lemma 2.2.** *Let  $q$  be a positive integer, and let  $p > 2q$  be a prime. Then, for any integer  $k$  with  $q-1 \leq k \leq p-1$  we have*

$$\sum_{q \leq j \leq k} \frac{1}{j^2} + \sum_{q \leq j < p+q-1-k} \frac{1}{j^2} \equiv \sum_{j=q}^{p-q} \frac{1}{j^2} + \sum_{0 \leq l < q-1} \frac{1}{(k-l)^2} \pmod{p}. \quad (2.1)$$

*Proof.* Observe that

$$\begin{aligned} \sum_{q \leq j < p+q-1-k} \frac{1}{j^2} &\equiv \sum_{q \leq j < p+q-1-k} \frac{1}{(p-j)^2} = \sum_{k+1-q < i \leq p-q} \frac{1}{i^2} \\ &= \sum_{0 < i < q} \frac{1}{i^2} + \sum_{i=q}^{p-q} \frac{1}{i^2} - \sum_{0 < i \leq k+1-q} \frac{1}{i^2} \pmod{p} \end{aligned}$$

and

$$\sum_{q \leq j \leq k} \frac{1}{j^2} + \sum_{0 < i < q} \frac{1}{i^2} - \sum_{0 < i \leq k+1-q} \frac{1}{i^2} = \sum_{k+1-q < j \leq k} \frac{1}{j^2} = \sum_{0 \leq l < q-1} \frac{1}{(k-l)^2}.$$

So the desired (2.1) follows.  $\square$

**Lemma 2.3.** *Let  $m > 2$  and  $q > 0$  be integers. Then, for any prime  $p > mq$  we have*

$$\sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \equiv 0 \pmod{p} \quad (2.2)$$

and

$$\sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{0 \leq l < q-1} \frac{1}{(k-l)^2} \equiv 0 \pmod{p}. \quad (2.3)$$

*Proof.* As  $m(q-1) \leq mq-1 \leq p-2$ , we can write

$$\binom{x}{q-1}^m = \sum_{j=0}^{p-2} a_j x^j$$

with  $a_0, \dots, a_{p-2}$   $p$ -adic integers. It is well-known that  $\sum_{k=0}^{p-1} k^j \equiv 0 \pmod{p}$  for any positive integer  $j \not\equiv 0 \pmod{p-1}$  (see, e.g., [3, p. 235]). Thus

$$\sum_{k=q-1}^{p-1} \binom{k}{q-1}^m = \sum_{k=0}^{p-1} \sum_{j=0}^{p-2} a_j k^j = \sum_{j=0}^{p-2} a_j \sum_{k=0}^{p-1} k^j \equiv 0 \pmod{p}.$$

Let  $l$  be any integer with  $0 \leq l < q-1$ . Then

$$\begin{aligned} & \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \frac{1}{(k-l)^2} \\ & \equiv \sum_{k=l+1}^{p+l-1} \binom{k}{q-1}^m \frac{1}{(k-l)^2} = \sum_{k=1}^{p-1} \binom{k+l}{q-1}^m \frac{1}{k^2} \pmod{p}. \end{aligned}$$

As  $m(q-1) < mq \leq p-1$  and  $\binom{l}{q-1} = 0$ , we may write

$$\binom{x+l}{q-1}^m = \sum_{m \leq j < p-1} c_j x^j$$

with  $c_m, \dots, c_{p-2}$   $p$ -adic integers. Hence

$$\begin{aligned} \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \frac{1}{(k-l)^2} & \equiv \sum_{k=0}^{p-1} \sum_{m \leq j < p-1} c_j k^{j-2} \\ & \equiv \sum_{m \leq j < p-1} c_j \sum_{k=0}^{p-1} k^{j-2} \equiv 0 \pmod{p}. \end{aligned}$$

Now it follows from the above that

$$\begin{aligned} & \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{0 \leq l < q-1} \frac{1}{(k-l)^2} \\ & = \sum_{0 \leq l < q-1} \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \frac{1}{(k-l)^2} \equiv 0 \pmod{p}. \end{aligned}$$

This concludes the proof.  $\square$

## 3. PROOF OF THEOREM 1.2

*Proof of Theorem 1.2.* For each  $k \in \{1, 2, \dots, p-1\}$ , obviously

$$(-1)^{km} \binom{p/m - q}{k}^m = (-1)^{km} \prod_{j=1}^k \left( \frac{p/m - q - j + 1}{j} \right)^m = \prod_{j=1}^k \left( 1 + \frac{q-1}{j} - \frac{p}{jm} \right)^m$$

is congruent to

$$\begin{aligned} & \prod_{j=1}^k \left( \left( 1 + \frac{q-1}{j} \right)^m - \left( 1 + \frac{q-1}{j} \right)^{m-1} \frac{pm}{jm} + \frac{m(m-1)}{2} \left( 1 + \frac{q-1}{j} \right)^{m-2} \frac{p^2}{j^2 m^2} \right) \\ &= \prod_{j=1}^k \left( \left( 1 + \frac{q-1}{j} \right)^m - \left( 1 + \frac{q-1}{j} \right)^{m-1} \frac{p}{j} + \frac{m-1}{2m} \left( 1 + \frac{q-1}{j} \right)^{m-2} \frac{p^2}{j^2} \right) \end{aligned}$$

module  $p^3$ , and hence

$$\begin{aligned} & (-1)^{km} \binom{p/m - q}{k}^m - \prod_{j=1}^k \left( \left( 1 + \frac{q-1}{j} \right)^m - \left( 1 + \frac{q-1}{j} \right)^{m-1} \frac{p}{j} \right) \\ &\equiv \frac{m-1}{2m} \sum_{j=1}^k \left( 1 + \frac{q-1}{j} \right)^{m-2} \frac{p^2}{j^2} \prod_{\substack{i=1 \\ i \neq j}}^k \left( 1 + \frac{q-1}{i} \right)^m \\ &= p^2 \frac{m-1}{2m} \sum_{j=1}^k \frac{1}{(j+q-1)^2} \prod_{i=1}^k \left( 1 + \frac{q-1}{i} \right)^m \\ &= p^2 \frac{m-1}{2m} \binom{k+q-1}{k}^m \sum_{j=1}^k \frac{1}{(j+q-1)^2} \pmod{p^3}. \end{aligned}$$

Note that

$$\begin{aligned} & \prod_{j=1}^k \left( \left( 1 + \frac{q-1}{j} \right)^m - \left( 1 + \frac{q-1}{j} \right)^{m-1} \frac{p}{j} \right) \\ &= \prod_{j=1}^k \left( 1 + \frac{q-1}{j} \right)^{m-1} \prod_{j=1}^k \left( 1 + \frac{q-1}{j} - \frac{p}{j} \right) \\ &= \binom{k+q-1}{k}^{m-1} (-1)^k \binom{p-q}{k}. \end{aligned}$$

So, from the above we obtain

$$\begin{aligned}
& \sum_{k=0}^{p-1} (-1)^{km} \binom{p/m - q}{k}^m - \sum_{k=0}^{p-1} \binom{k+q-1}{k}^{m-1} (-1)^k \binom{p-q}{k} \\
& \equiv \sum_{k=0}^{p-1} p^2 \frac{m-1}{2m} \binom{k+q-1}{q-1}^m \sum_{0 < j \leq k} \frac{1}{(j+q-1)^2} \\
& = p^2 \frac{m-1}{2m} \sum_{k=q-1}^{p+q-2} \binom{k}{q-1}^m \sum_{q \leq j \leq k} \frac{1}{j^2} \\
& \equiv p^2 \frac{m-1}{2m} \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{q \leq j \leq k} \frac{1}{j^2} \pmod{p^3}.
\end{aligned}$$

Clearly,

$$\begin{aligned}
& 2 \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{q \leq j \leq k} \frac{1}{j^2} \\
& = \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{q \leq j \leq k} \frac{1}{j^2} + \sum_{k=q-1}^{p-1} \binom{p+q-2-k}{q-1}^m \sum_{q \leq j < p+q-1-k} \frac{1}{j^2},
\end{aligned}$$

and

$$\binom{p+q-2-k}{q-1}^m \equiv \binom{q-2-k}{q-1}^m = (-1)^{m(q-1)} \binom{k}{q-1}^m = \binom{k}{q-1}^m \pmod{p}$$

for all  $k = q-1, q, \dots, p-1$ . Therefore, with the helps of Lemmas 2.2 and 2.3, we get

$$\begin{aligned}
& 2 \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \sum_{q \leq j \leq k} \frac{1}{j^2} \\
& \equiv \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \left( \sum_{q \leq j \leq k} \frac{1}{j^2} + \sum_{q \leq j < p+q-1-k} \frac{1}{j^2} \right) \\
& \equiv \sum_{k=q-1}^{p-1} \binom{k}{q-1}^m \left( \sum_{j=q}^{p-q} \frac{1}{j^2} + \sum_{0 \leq l < q-1} \frac{1}{(k-l)^2} \right) \\
& \equiv 0 \pmod{p}.
\end{aligned}$$

As the degree of the polynomial  $\binom{x+q-1}{q-1}^{m-1}$  is  $(m-1)(q-1) < (m-1)q < p-q$ , by Lemma 2.1 we have

$$\sum_{k=0}^{p-1} \binom{k+q-1}{k}^{m-1} (-1)^k \binom{p-q}{k} = \sum_{k=0}^{p-q} \binom{p-q}{k} (-1)^k \binom{k+q-1}{q-1}^m = 0.$$

Therefore the desired (1.1) follows from the above.  $\square$

## REFERENCES

- [1] V.J.W. Guo and J. Zeng, *Some  $q$ -analogues of supercongruences of Rodriguez-Villegas*, J. Number Theory **145** (2014), 301–316.
- [2] C. Helou, G. Terjanian, *On Wolstenholme’s theorem and its converse*, J. Number Theory **128** (2008), 475–499.
- [3] K. Ireland, M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd Edition, Grad. Texts in Math., Vol. 84, Springer, New York, 1990.
- [4] Z.-W. Sun, *Supercongruences involving products of two binomial coefficients*, Finite Fields Appl. **22** (2013), 24–44.
- [5] Z.-W. Sun, *Supercongruences motivated by  $e$* , J. Number Theory **147** (2015), 326–341.
- [6] J.H. van Lint, R.M. Wilson, *A Course in Combinatorics*, 2nd Edition, Cambridge Univ. Press, Cambridge, 2011.
- [7] J. Wolstenholme, *On certain propertities of prime numbers*, Quart. J. Appl. Math. **5** (1862), 35–39.

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, PEOPLE’S REPUBLIC OF CHINA

*E-mail address:* xzmeng@smail.nju.edu.cn and zwsun@nju.edu.cn